

specimen was identified by Adrian Riley and is retained at the Yarner Wood Forest Office. As far as I am aware this was the first specimen of this species to be captured in Devon.

A specimen of the Sword Grass *Xylena exsoleta* L. (Noctuidae) was captured in the Rothamsted trap in Yarner Wood, Dartmoor, Devon on 7.xii.1994. The specimen is retained at the Yarner Wood Forest Office and was subsequently confirmed by myself on 25.vii.1996. As far as I can see, there have been seven other historic sightings of this species in Devon.

A specimen of Haworth's Minor *Celaena haworthii* Curt. was taken by Mr A. Jenkins at Chardstock, Devon on 18.viii.1996; there have been specimens of this species taken on Dartmoor in the past although none of these records have been published. If anyone can help to rectify this lack of records of this species it would prove to be very useful. I would like to thank Mr Jenkins for allowing me to publish this finding; the specimen is in the collection of the author of this note.

Finally, following my earlier note (*Ent. Rec.* **108**: 148), I have now confirmed that the Bloxworth Snout *Hypena obsitalis* Hb. is in fact breeding at Churston, South Devon. Larvae were collected on 2.vii.1996 and were bred through to adult on 18.vii.1996 (male) and 22.vii.1996 (female).

– ROY MCCORMICK, Devonshire Lepidoptera Recorder, 36 Paradise Road, Teignmouth, Devon TQ14 8NR.

Observation of a second generation Orange-tip *Anthocharis cardamines* L. (Lep.: Pieridae)

Throughout most of Hertfordshire, the Orange-tip remains one of our most common butterflies. The 1996 season proved no exception, with large numbers being observed on the Environmental Change Network (ECN) common butterfly census conducted on the Rothamsted estate. Males of the Orange-tip are usually on the wing in late April and early May. The earliest siting for Hertfordshire is 13 April 1961 and the latest, 8 July 1902 (Sawford, 1987, *The Butterflies of Hertfordshire*, Castlemead, Ware). Very occasionally, a second generation may emerge in late summer, (Sawford, 1987 *op. cit.*, Thomas and Lewington, 1991, *The Butterflies of Britain and Ireland*, Dorling Kindersley, London). On 5 September 1996, on a sunny afternoon with a light breeze, whilst working in a garden in the village of Redbourn, OS grid reference TL 103122, a male Orange-tip flew in and remained briefly but long enough to observe its characteristic orange wing-tips. This observation is nearly two months later than any previous record for Hertfordshire. – JOHN E. BATER, Department of Entomology and Nematology, IACT Rothamsted, Harpenden, Hertfordshire AL5 2JQ.